

Homecare After Dental Surgery - Dental Implant Surgery

After dental surgery or dental implant surgery, it's important to follow these recommendations to ensure their long term success.

General Care

To control minor bleeding, apply moderate pressure with gauze pads or a clean moist tea bag wrapped in gauze.

Limit yourself to calm activities for the first three to five days to keep your blood pressure stable and reduce bleeding.

If you usually wear a removable denture or partial denture, we will give you specific instructions about wearing it while you are healing.

Tobacco Use

Don't use tobacco for at least 24 hours, and avoid tobacco as much as possible for several months. Tobacco slows healing and could cause a very painful dry socket.

When to call us

Call us right away if you have heavy or increased bleeding, you have pain or swelling that increases or continues beyond two or three days, or you have a reaction to the medication.

Call us if the fragments gradually show through the gums, so we can check them.

You can also call us with any questions or concerns.

Chewing and Eating

Avoid chewing until the numbness has worn off completely.

For the first 24 hours, drink lots of liquids and eat soft, nutritious foods, chewing on the opposite side of your mouth.

After the first 24 hours, you can begin to return to your usual diet, but continue to chew on the opposite side. Avoid very hot or spicy foods until your gums are healed.

Brushing and Flossing

For the first 14 days, gently wipe the surgery site with gauze or a cotton swab. After that, brush the area gently with a soft toothbrush.

Don't floss the site for four weeks. We will evaluate the healing and adjust your brushing and flossing timetable as necessary.

Brush and floss your other teeth normally.

If we have recommended a mouth rinse, use it as directed.

Medication and Discomfort

If antibiotics or other medicines are prescribed, continue to take them for the indicated length of time, even if all symptoms and signs of infection are gone.

To control discomfort, take pain medication before the anesthetic has worn off or as recommended. It's normal to experience some discomfort for several days after surgery.

To reduce discomfort or swelling for the first 24 hours following surgery use ice packs, 20 minutes on and 20 minutes off.

To reduce discomfort after the first 24 hours, apply moist heat.

To further reduce discomfort **after** the first 24 hours, you can rinse your mouth very gently three times a day with warm salt water. Use about one teaspoon of salt per glass of warm water.

Post-Operative Pain Control

The following protocol will be of benefit to patients, provided they are not allergic to aspirin, Tylenol or Ibuprofen.

Take 600 mg of Ibuprofen (trade names: Advil, Nuprin, Motrin IB, Or just generic Ibuprofen 200 mg). To get a total of 600 mg, you must take 3 tablets of the 200 mg form. We prefer 600mg (3 tablets) to 200 mg or 400 mg because the first 400 mg is for pain relief, the next 200 mg is an anti-inflammatory in nature and this is what actually helps prevent pain. If in the event the 600 mg ibuprofen upsets your stomach, cut back to 1 or 2 tablets. Take every four to six hours.

If the Ibuprofen does not control your pain, then alternate taking 2 tablets of Tylenol Extra Strength (Tylenol 500 mg, generic acetaminophen 500 mg) every four to six hours with the ibuprofen 600 mg.

If you have any questions, call the office, 216-524-4410.

